

- Hoerstrup, Zund, Lachat, Schoeberlein, Uhlschmid, Vogt, Turina, "Tissue engineering: a new approach in cardiovascular surgery-seeding of human fibroblasts on resorbable mesh," *Swiss Surgery*, (Suppl.), 2:23-5, 1998.
- Hoerstrup, Zund, Schoeberlein, Ye, Vogt, Turina, "Fluorescence activated cell sorting: a reliable method in tissue engineering of a bioprosthetic heart valve," *Annals of Thoracic Surgery*, 66(5):1653-7, 1998.
- Isogai, Landis, Kim, Gerstenfeld, Upton, Vacanti, "Formation of phalanges and small joints by tissue-engineering," *J. of Bone & Joint Surgery, American Vol.*, 81(3):306-16, 1999.
- Martin, Padera, Vunjak-Novakovic, Freed, "In vitro differentiation of chick embryo bone marrow stromal cells into cartilaginous and bone-like tissues," *J. of Orthopaedic Res.*, 16(2):181-9, 1998.
- Nagy et al., "Pathogenesis of ascites tumor growth: vascular permeability factor, vascular hyperpermeability, and ascites fluid accumulation," *Cancer Res.*, 55:360, 1995.
- Peppas and Langer, "New challenges in biomaterials," *Science*, 263:1715-1720, 1994.
- Peter, Miller, Yasko, Yaszemski, Mikos, "Polymer concepts in tissue engineering," *J. of Biomed. Materials Res.*, 43(4):422-7, 1998.
- Sacks, Chuong, Petroll, Kwan, Halberstadt, "Collagen fiber architecture of a cultured dermal tissue," *J. of Biomed. Engineering*, 119(1):124-7, 1997.
- Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989.
- Shinoka, Shum-Tim, Ma, Tanel, Isogai, Langer, Vacanti, Mayer, "Creation of viable pulmonary artery autografts through tissue engineering," *J. of Thoracic & Cardiovascular Surgery*, 115(3):536-45, 1998.
- Sims, Butler, Cao, Casanova, Randolph, Black, Vacanti, Yaremchuk, "Tissue engineered neocartilage using plasma derived polymer substrates and chondrocytes," *Plastic & Reconstructive Surgery*, 101(6):1580-5, 1998.
- Vunjak-Novakovic, Obradovic, Martin, Bursac, Langer, Freed, "Dynamic cell seeding of polymer scaffolds for cartilage tissue engineering," *Biotechnology Progress*, 14(2):193-202, 1998.
- Whang, Tsai, Nam, Aitken, Sprague, Patel, Healy, "Ectopic bone formation via rhBMP-2 delivery from porous bioabsorbable polymer scaffolds," *J. of Biomed. Materials Res.*, 42(4):491-9, 1998.
- Wong and Mooney, "Synthesis and properties of biodegradable polymers used in tissue engineering," In: *Synthetic Biodegradable Polymer Scaffolds*, (Atala and Mooney, eds.), Birkhauser Press, Boston, Mass., pp. 51-82, 1997.
- Yoo and Atala, "A novel gene delivery system using urothelial tissue engineered neorgans," *J. of Urology*, 158(3 Pt 2):1066-70, 1997.
- What is claimed is:

1. A composition comprising at least one biodegradable polymer fiber wherein said fiber is composed of a first phase and a second phase, the first and second phases being immiscible, and wherein the second phase comprises one or more therapeutic agents.
2. The composition of claim 1, wherein said second phase is derived from an aqueous solution, a hydrogel or polymer.
3. The composition of claim 1, wherein said fiber forms a scaffold and further wherein, said second phase is manipulated to form an internal porous structure within the fiber.
4. The composition of claim 1, wherein said fiber is woven, braided or knitted in an assembly with other fibers, and at least one fiber in the assembly comprises one or more therapeutic agents.

5. The composition of claim 1, wherein the one or more therapeutic agents are distributed within the second phase in a nonhomogenous pattern.
6. The composition of claim 1, wherein the concentration of said one or more therapeutic agents varies along the longitudinal axis of the fiber.
7. The composition of claim 6, wherein the concentration of said one or more therapeutic agents varies linearly, exponentially or in any desired fashion, as a function of distance along the longitudinal axis of the fiber.
8. The composition of claim 1, wherein the concentration of said one or more therapeutic agents decreases from a first end of the fiber to a second end of the fiber.
9. The composition of claim 1, wherein said one or more therapeutic agents vary in a bidirectional manner, and the content of said one or more therapeutic agents increases from the first end of said fiber to a maximum and then decreases towards the second end of said fiber.
10. The composition of claim 1, further comprising at least one biodegradable polymer fiber containing no therapeutic agent.
11. The composition of claim 1, wherein said one or more therapeutic agents are selected from the group consisting of drugs, proteins, enzymes, growth factors, immunomodulators, compounds promoting angiogenesis, compounds inhibiting angiogenesis, anti-inflammatory compounds, antibiotics, cytokines, anti-coagulation agents, procoagulation agents, chemotactic agents, agents to promote apoptosis, agents to inhibit apoptosis, and mitogenic agents.
12. The composition of claim 1, wherein said one or more therapeutic agents include a radioactive agent or a contrast agent for imaging studies.
13. The composition of claim 1, wherein said one or more therapeutic agents is selected from the group consisting of viral vector, polynucleotide and polypeptide.
14. The composition of claim 1, wherein said one or more therapeutic agents comprise an angiogenesis-promoting agent.
15. The composition of claim 14, wherein said angiogenesis-promoting agent is vascular endothelial growth factor.
16. The composition of claim 1, wherein said biodegradable polymer is a single polymer, a co-polymer, or a mixture of polymers selected from the group consisting of polypeptides, polydepsipeptides, nylon copolyamides, aliphatic polyesters, polydihydropyrans, polyphosphazenes, poly(ortho ester), poly(cyano acrylates), polyanhydride, modified polysaccharides and modified proteins.
17. The composition of claim 16, wherein said aliphatic polyesters are selected from the group consisting of poly(glycolic acid), poly(lactic acid), poly(alkylene succinates) poly(hydroxybutyrate), poly(butylene diglycolate), poly(epsilon-caprolactone) and copolymers, blends and mixtures thereof.
18. The composition of claim 16, wherein said modified polysaccharides are selected from the group consisting of cellulose, starch-alginate and the glycosaminoglycans, chondroitin sulfate, heparin, heparin sulfate, dextran, dextran sulfate, chitin, chitosan and chitosan sulfate.
19. The composition of claim 16, wherein said modified proteins are selected from the group consisting of collagen and fibrin.
20. The composition of claim 1, wherein said fiber comprises a plurality of polymer layers, wherein an outer layer circumscribes an adjacent inner layer.
21. The composition of claim 20, wherein said plurality of layers optionally contain one or more therapeutic agents.